

$\mathcal{N} = 1$ Heterotic String Vacua from Mirror Symmetry ¹

P. Mayr

Theory Division, CERN, 1211 Geneva 23, Switzerland

Abstract

We review a systematic construction of $\mathcal{N} = 1$ supersymmetric heterotic string vacua using mirror symmetry. The method provides a large class of explicit solutions for stable, holomorphic vector bundles on Calabi–Yau n -folds Z_n in terms of toric geometry. Phenomenologically interesting compactifications as well as non-perturbative dynamics of the heterotic string are discussed within this framework.

April 1999

¹ Lectures presented at the Winter School on Vector Bundles, Mirror Symmetry, and Lagrangian Submanifolds, Harvard University, March 1999

1. Introduction

Heterotic string compactifications with minimal $\mathcal{N} = 1$ supersymmetry have been the phenomenologically most promising candidate for a unifying theory of particle physics and gravitation for a while. In addition to having the smallest possible amount of supersymmetry, which is supposed to provide, amongst other, a solution to the problem of why there is such a huge discrepancy between the scales determining the gravitational and gauge interactions, these theories have rich gauge symmetries, namely $E_8 \times E_8$ or $SO(32)$, in ten dimensions. Those should be broken to the gauge groups of standard particle physics, be it grand unified groups or that of the standard model.

The mathematical conditions that ensure $\mathcal{N} = 1$ supersymmetry in a compactification on a compact manifold times flat d -dimensional Minkowski space have been determined a long time ago [1][2][3]. However explicit solutions to these equations are rather hard to obtain. In this lecture we review a geometric construction of a large class of such heterotic vacua using a generalized version of mirror symmetry in type II strings. In particular we define a map $f : \mathcal{W}_{n+1} \rightarrow (Z_n, V)$ that associates to a (toric) local $n + 1$ dimensional Calabi–Yau geometry $\mathcal{W}_{n+1}$ the data of a (toric) Calabi–Yau n -fold Z_n together with a family of vector bundles V (or more generally sheaves) on it. The relation between the type II string compactifications and heterotic strings can be traced back to string dualities, namely type IIA/heterotic [4][5][6] and F-theory/heterotic duality [7][8]. Within this context we establish this duality at the classic level and provide a systematic construction of dual pairs $W_{n+1} \leftrightarrow (Z_n, V)$, where W_{n+1} is now the global $n + 1$ -fold used for F-theory compactification.

2. Heterotic $\mathcal{N} = 1$ vacua

The conditions for supersymmetric compactifications of perturbative heterotic strings on a manifold Z have been determined in [1][2][3] under some simplifying restrictions, such as the vanishing of the three-form field strength $H = dB + \dots$ and a constant dilaton background. These conditions are modified in non-perturbative vacua by the presence of magnetic five-branes [9].

2.1. Perturbative vacua

Supersymmetry in the uncompactified space-time requires the existence of a covariantly constant spinor on Z which means that Z is a Calabi–Yau manifold. In addition to the geometry one has to specify a gauge background V on Z with the following properties.

The vector bundle V has to satisfy the topological conditions

$$\begin{aligned} c_1(V) &= 0 \bmod 2, \\ \lambda(V) &= c_2(Z), \end{aligned} \tag{2.1}$$

where λ is the four-dimensional characteristic class of V . The first equation ensures that the bundle V admits spinors. The second equation derives from the Bianchi identity for the three-form field strength H .

Moreover supersymmetry requires that the connection on V satisfies

$$F_{ab} = F_{\bar{a}\bar{b}} = 0, \quad g_{a\bar{b}} F^{a\bar{b}} = 0. \tag{2.2}$$

So V is holomorphic and moreover the second equation says that V is a stable bundle.

The third Chern class of V is also important: it determines the net number of generations of the compactification as [3]

$$N_{gen} - N_{anti-gen} = \frac{1}{2}c_3(V). \quad (2.3)$$

Later it was realized that the relevant gauge backgrounds for string theory are not actually smooth vector bundles but rather include also more singular configurations such as reflexive or coherent sheaves [10]. These are also the objects that appear naturally in the geometric construction discussed in the following.

2.2. Non-perturbative vacua: five-branes

The understanding of non-perturbative dynamics of the heterotic string improved dramatically with the uprise of string dualities. The basic non-perturbative state of the heterotic theory is the magnetic dual of the fundamental string, the five-brane. In particular, adding background five-branes to a heterotic compactification can yield a new, non-perturbative compactification with the same maximal amount of supersymmetry as dictated by the compactification geometry [9]. Non-perturbative five-branes add an extremely rich spectrum of dynamics to the perturbative heterotic string, including non-critical strings [11] and non-abelian gauge symmetries of very high rank [12].

The presence of non-perturbative five-branes changes the topological constraint on the gauge background V . The magnetic five-brane contributes a source term to the Bianchi identity for the three-form H :

$$dH = tr R \wedge R - tr F \wedge F - q_{5B} \sum_{five-branes} \delta_{5B}^{(4)}, \quad (2.4)$$

which integrates on a four-cycle to

$$c_2(Z) = c_2(V) + [W]. \quad (2.5)$$

Here $\delta_{5B}^{(4)}$ is a formal four-form that integrates to one in the directions *transverse* to a single five-brane and $[W]$ denotes the class associated to this last term in (2.4). In other words, a five-brane is able to correct the mismatch of gravitational and gauge curvatures in the direction transverse to the world volume. Thus adding five-branes, which take care of part of the anomaly related to $c_2(Z)$, enlarges substantially the class of consistent gauge backgrounds on Z [13]. This interplay between gauge fields and five-branes is clear from the interpretation of the latter as the zero size limit of gauge instantons [14][11].

Lorentz invariant compactifications with five-branes are possible in $d \leq 6$ compactifications. In six dimensions, the curvature is localized along K3 and the five-branes simply fill space-time. These five-branes contribute also to the massless spectrum. On the five-brane of the $E_8 \times E_8$ string, there is an antisymmetric tensor B' with anti-self-dual field strength. It couples to anti-self-dual strings in 6d with a string tension parametrized by a real scalar ϕ' . There are four more real scalars ϕ_i which parametrize the position of the five-brane in the transverse space. There is also an $SO(5)_R$ R-symmetry under which ϕ', ϕ_i transform as a $\underline{5}$ and B' as a singlet.

In four uncompactified dimensions the five-brane world volume has to wrap a supersymmetric 2-cycle C_2 in Z , that is a holomorphic curve [15][16]. The massless spectrum depends now on the geometry of the 2-cycle C_2 . The answer can be obtained from the twisting of the world volume theory [16]. It would be interesting to obtain the general answer. Here we restrict to some simple interesting situations. A special

supersymmetric twisting for a five-brane compactified on a curve $C_2^{(g)}$ of genus g has been considered in [17] with the result that out of the five scalars, two become 1-forms on $C_2^{(g)}$ while three remain scalars $\tilde{\phi}_i$. In this case we get g vector fields from B' with one index in the $C_2^{(g)}$ direction, as well as $2g + 3$ real scalars and a single anti-symmetric tensor B' in four dimensions².

If $g = 1$, the normal bundle of $C_2^{(1)}$ is trivial and the above analysis applies. There are six scalars and a single vector field which combine to a $\mathcal{N} = 4$ vector multiplet. This situation is encountered if $C_2^{(1)}$ is the fiber of the elliptic fibration of Z . Enhancement to $SU(N)$ takes place if N five-branes coincide.

On the other hand consider the situation where C_2 is an isolated genus 0 curve in Z . Since B' is a singlet, it is unaffected by the twisting and must survive, similarly ϕ' which still parametrizes the tension of the four-dimensional string to which B' couples. Since C_2 is isolated there are no further scalars. The spectrum is that of an $\mathcal{N} = 1$ linear multiplet.

Apart from the generic spectrum for generic positions of the five-branes there will be additional light degrees of freedom for special values of the moduli. If several five-branes come close to each other, non-critical string with a vanishing tension will become relevant, similarly as in six dimensions. In the case of $g > 1$, winding states of the string contribute massless vector multiplets, charged with respect to the generic gauge fields from B' as in [17].

Another feature is the presence of extra matter in the perturbative heterotic gauge group for a special alignment of the gauge and geometric moduli [18]. This matter of non-perturbative origin is in addition to the contribution in (2.3).

Elliptically fibered Calabi–Yau manifolds Z

Generally, finding a suitable gauge background V fulfilling (2.1) and (2.2) explicitly is a very hard question, though there is an existence theorem for a solution in the generic case [19]. There is a much better understanding in the case where Z is elliptically fibered, basically because it is relatively simple to describe holomorphic (semi-)stable bundles on a torus E and fibering these data in a holomorphic way over a complex manifold B produces a holomorphic bundle V on the elliptic fibration $E \rightarrow Z \rightarrow B$. This situation has been considered in [13][20][21].

In the following we will take a quite different route to the subject of holomorphic stable vector bundles (or rather sheafs) on such n -dimensional Calabi–Yau manifolds Z_n . We define them purely in terms of type IIA string theory. The fact that the non-abelian gauge symmetries of type II arise from D-brane geometries will lead us to a unified (toric) geometric description of bundles on Z_n in terms of type IIA compactification geometries.

² Note that it was the non-compactness of C_2 in [17] that kills the scalars from 0-forms on $C_2^{(g)}$.

3. Vector bundles and type II strings

Instead of the heterotic string, we will consider now the ten-dimensional type IIA string compactified on Calabi–Yau $n + 1$ -fold singularities. Though it is not obvious at first sight, this theory has the same moduli space $\mathcal{M}_V$ of holomorphic stable bundles on a (different) Calabi–Yau manifold Z_n .

We will derive this statement in two steps, for $n = 2$ and $n > 2$. For $n = 2$ we have a relation between Wilson lines on a T^2 and Calabi–Yau 2-fold singularities which follows from a symmetry of the type IIA compactification on $K3 \times T^2$ that is closely related to mirror symmetry of K3. In the second step we use the adiabatic principle [22] and consider holomorphic fibrations of the 2-fold singularities to construct Calabi–Yau $n + 1$ -fold singularities related to holomorphic stable bundles on Calabi–Yau Z_n .

3.1. Mirror symmetry on K3 and generalizations

Mirror symmetry of K3

Consider a type IIA string compactified on a K3 surface. This gives an $\mathcal{N} = 2$ supersymmetric theory in six dimensions. From dimensional reduction we obtain twenty $\mathcal{N} = 2$ vector multiplets which contain as their bosonic degree of freedom each one vector and four real scalars. The latter transform as a $\underline{4}$ under an $SO(4)_R$ R-symmetry and parametrize the 58 metric moduli of K3 plus 22 values of the B -fields on $H^2(K3)$.

In a given algebraic realization M_2 of K3, the 80 moduli of K3 split in complexified Kähler and complex structure moduli³. The $SO(4)_R$ rotations acting on the scalars in $\underline{4}$ do not preserve this split in general. In particular there is a $SO(4)_R$ rotation that exchanges Kähler and complex structure moduli spaces $\mathcal{M}_{KM}$ and $\mathcal{M}_{CS}$, respectively. Such a transformation is usually known as *mirror symmetry* of K3 manifolds, which says that a type IIA string compactified on M_2 describes the same physics as a type IIA string compactification on a "different"⁴ K3 manifold W_2 . The new algebraic K3 surface W_2 is called the mirror manifold of M_2 . We have the following identifications under mirror symmetry:

$$\begin{aligned} M_2 &\leftrightarrow W_2, \\ \mathcal{M}_{CS}(M_2) &\leftrightarrow \mathcal{M}_{KM}(W_2), \\ \mathcal{M}_{KM}(M_2) &\leftrightarrow \mathcal{M}_{CS}(W_2). \end{aligned} \tag{3.1}$$

An extension of mirror symmetry

There is an intriguing generalization of this symmetry upon compactification on a further T^2 to four dimensions. In this case we obtain a $\mathcal{N} = 4$ supersymmetric string theory. The $\mathcal{N} = 4$ vector multiplet contains now 6 scalars which transform in a $\underline{6}$ of an $SO(6)_R$ symmetry. The two extra scalars are the internal components A_i of the six-dimensional gauge fields on the T^2 , call it E , and thus parametrize the stringy moduli space of Wilson lines on the elliptic curve, $\mathcal{M}_E^{string}(H)$. Here H denotes the structure group of the Wilson line background. Again the $SO(6)_R$ rotations provide identifications within the type IIA moduli space $\mathcal{M}_{IIA}(K3 \times T^2)$:

$$\mathcal{M}_{KM}(M_2) \leftrightarrow \mathcal{M}_{CS}(W_2) \leftrightarrow \mathcal{M}_E^{string}(H). \tag{3.2}$$

³ Subscripts, as in M_2 , denote the complex dimension of a geometry.

⁴ Since the K3 surface is unique, mirror symmetry is actually a discrete identification in the moduli space of K3 surfaces.

In particular there is now an element of $SO(6)_R$, corresponding to the last arrow, which provides a relation between the moduli space of stringy Wilson lines with the Kähler moduli space of an algebraic K3 manifold.

The dual heterotic picture

Let us illustrate the symmetry (3.2) of the type IIA compactification on $K3 \times T^2$ in the dual heterotic picture where it is very obvious; however we stress that we do not need the heterotic dual to derive the symmetry. Fig. 1 shows the type IIA compactification and the corresponding dual heterotic string compactified on $T_1^2 \times T_2^2 \times T_3^2$. The factorization of T^6 is related to *i*) the elliptic fibration and *ii*) the factorization of the gauge background on $K3 \times T^2$ in the type IIA theory. Under these circumstances, the Kähler moduli of K3 can be identified with moduli space $\mathcal{M}_{T_1^2}(H_1)$ of H_1 Wilson lines on the first T_1^2 and similarly complex structure moduli with Wilson lines in the second factor $\mathcal{M}_{T_2^2}(H_2)$. The non-abelian gauge symmetry in six dimensions is G which on the type IIA side arises from the G singularity and on the heterotic side is the commutant of $H_1 \times H_2$ in G_0^{het} . Upon further compactification to four dimensions on a torus T_3^2 , we obtain another factor $\mathcal{M}_{T_3^2}(G)$ on both sides.




Fig. 1: The symmetry (3.2) from a heterotic point of view.

The heterotic theory has an obvious symmetry under permutation of the three T^2 factors. The exchange of the first two tori amounts to mirror symmetry in the type IIA theory. On the other hand an exchange that involves the third torus gives a new symmetry of the moduli space of the type IIA compactification that identifies Wilson lines with structure group G on T^2 with either Kähler or complex deformations of the K3. This is the new relation described in (3.2).

3.2. The precise correspondence

We want now to use the equivalence (3.2) implied by the $SO(6)_R$ transformations to describe the moduli space $\mathcal{M}_E(H)$ of flat holomorphic (and non-abelian) H bundles on an elliptic curve in terms of complex deformations of a geometric type IIA compactification. There are a few basic considerations to make precise such a correspondence. We will not dive into the details here, which can be found in [23][18], but rather sketch the procedure and state the result.

- Firstly note that the field theoretical moduli space $\mathcal{M}_E(H)$ contains also the deformations of the torus E . This is related to the fact that we have to consider gauge backgrounds for the $K3 \times T^2$ compactification which factorize in the sense that they do not restrict the geometry of the T^2 factor.
- The moduli space $\mathcal{M}_E^{string}(H)$ that appears in (3.2) is still that of a theory of Wilson lines coupled to *string* theory. To obtain the field theoretical moduli space $\mathcal{M}_E$ we have to decouple the string states by sending the string scale to infinity, $M_{str} \rightarrow \infty$.

- c) The structure group H of the Wilson lines on T^2 is contained in the gauge group of the six-dimensional compactification on M_2 , which is generically abelian. Non-abelian gauge symmetries in six dimensions arise from charged vector multiplets associated to D2 brane wrappings on 2-cycles C^i in M_2 [5]. The masses of these states are $\sim M_{str} \cdot Vol(C^i)$. Therefore light non-abelian vector multiplets are associated to very small 2-cycles in M_2 in the limit $M_{str} \rightarrow \infty$.

The condition *a)* translates to the property that M_2 has to be elliptically fibered. Conditions *b)* and *c)* are well-known from the geometric engineering of $\mathcal{N} = 2$ quantum field theories [24]. Taking the string decoupling limit and keeping at the same time the masses of the non-abelian states finite results in a geometric limit of K3 which is captured by a *local singularity* of the K3. Singularities of K3 at finite distance in the moduli space are the well-known ADE singularities. Their local versions are described in terms of ADE singularities of ALE spaces.

The local geometry is then of the form of an elliptic fibration over the $\mathbf{C}$ plane with an H singularity, with $H \subset ADE$, at the origin. There are $r = \text{rank } H$ 2-spheres C^i associated to the blow up of this singularity which intersect according to an ordinary H Dynkin diagram. The elliptic fibration implies the existence of a further 2-sphere C^0 associated to the generic homology class of the elliptic fiber; the intersections with the r 2-cycles C^i are described by the *affine* Dynkin diagram of H . In summary we arrive at the following equivalence [23][18]:

- o Let $\mathcal{M}_2$ denote the local neighbourhood of an H singularity in an elliptic fibration over the plane. There are $r+1$ volumes associated to C^k , $k = 0, \dots, r$ which parametrize the Kähler moduli space $\mathcal{M}_{KM}(\mathcal{M}_2)$ of $\mathcal{M}_2$. The type IIA compactification on $\mathcal{M}_2$ describes holomorphic stable H bundles on E with a moduli space parametrized by the $r+1$ Kähler moduli of $\mathcal{M}_2$:

$$M_{KM}(\mathcal{M}_2) \cong M_E(H). \quad (3.3)$$

Applying local mirror symmetry to the geometry $\mathcal{M}_2$ we obtain an equivalent description in terms of complex structure deformations of the mirror geometry $\mathcal{W}_2$:

- o Let $\mathcal{W}_2$ denote the (local) mirror of $\mathcal{M}_2$. The type IIA compactification on $\mathcal{W}_2$ describes holomorphic stable H bundles on E parametrized by the $r+1$ complex moduli of $\mathcal{M}_2$:

$$M_{CS}(\mathcal{W}_2) \cong M_E(H). \quad (3.4)$$

3.3. The complex geometries $\mathcal{W}_2$

Let us briefly summarize the structure of the two-complex dimensional geometries $\mathcal{W}_2$ which are the mirrors of the local ADE singularities in the elliptic fibration over the plane. We will recover in this way the results of [13] for those H where a geometric realization of $\mathcal{M}_E(H)$ has been given and reinterpret them in terms of type IIA compactifications. Moreover we obtain a unified and general geometric formulation for any H .

The geometry $\mathcal{W}_2$ is given as a hypersurface defined by the vanishing of a quasi-homogeneous polynomial in some variables (y, x, z, v) on a (toric) ambient space. Its general form is

$$p_{\mathcal{W}_2} = p_0(y, x, z) + p_+(y, x, z; v) = p_0 + \sum_{i=1} v^i p_+^i(y, x, z) = 0, \quad (3.5)$$

where we have separated the v independent term p_0 , and all polynomials are quasi-homogeneous in (y, x, z) . The v independent term p_0 is universal for all H and given by

$$p_0 = y^2 + x^3 + z^6 + \mu yxz. \quad (3.6)$$

$p_0 = 0$ describes an elliptic curve with complex structure modulus μ . The v dependent term depends on the structure group H and has an intriguing group theoretical structure: consider the affine Dynkin diagram $\Gamma(H)$ associated to the Lie group H . Then each node in $\Gamma(H)$ with Dynkin index s_i contributes one monomial to $p_+^i(y, x, z)$. See appendix A for an illustration of this fact.

Let us give one simple example. For $H = A_{N-1}$, $\Gamma(A_{N-1})$ contains N nodes with Dynkin index one. The equation defining $\mathcal{W}_2(A_{N-1})$ obtained from local mirror symmetry (for N even) is

$$p_0 + v (a_N z^N + a_{N-2} z^{N-2} x + a_{N-3} z^{N-3} y + \dots + a_0 x^{N/2}) = 0, \quad (3.7)$$

with p_0 as above. This defines a two complex dimensional local geometry with a complex structure determined by the complex parameters μ, a_k . In fact μ defines the complex structure of the torus $\hat{E} : p_0 = 0$ on which the bundle is defined. The a_k are coordinates on the $\mathbf{P}^{N-1}$ predicted in the mathematics literature [25]. It is well-known, that as far as complex structure is concerned, linear variables in the defining "superpotential" can be simply integrated out, which amounts to setting $dW/dv = 0$ in addition to $W = 0$. In our case, varying with respect to v gives $p_+ = 0$ and $p_0 = 0$ separately. This is the zero-dimensional spectral cover description found in [13].

Moreover, from the above construction, we get a similar description for all structure groups H in terms of complex deformations of a two-dimensional complex geometry $\mathcal{W}_2(H)$ which is identified as a physical type IIA compactification geometry.

Interestingly, we can even describe more sophisticated bundles, such as the $Spin(32)/\mathbf{Z}_2$ bundles without vector structure considered in [26], in terms of two-complex dimensional geometries $\mathcal{W}_2$. These bundles lead to a disconnected set of vacua for the heterotic $Spin(32)/\mathbf{Z}_2$ string, the CHL vacua [27] with maximal supersymmetry but reduced rank of the gauge group. The bundles associated to these vacua can be constructed in terms of special elliptic fibrations with reduced monodromy group of the fibration [28].

3.4. Vector bundles on Calabi–Yau n -folds

Apart from the satisfying formulation of holomorphic H bundles on a torus, unifying and extending the results obtained in [13], the real pay-off of the geometric description derives from the type IIA compactification picture. Combining the above construction with an adiabatic argument as in [29], we can get local type IIA compactifications that describe holomorphic stable vector bundles on Calabi–Yau n -folds in a simple way. All we have to do is to fiber the two-dimensional geometries $\mathcal{W}_2$ holomorphically over a $n - 1$ dimensional base B_{n-1} , such that the total geometry describes a local patch of a type IIA compactification on a Calabi–Yau $n + 1$ -fold $\mathcal{W}_{n+1}$. The general form of $\mathcal{W}_{n+1}$ is as in (3.5), with the difference that the complex parameters defining the complex structure, i.e. μ, a_k in the A_{N-1} case, now get upgraded to sections of line bundles on the base B_{n-1} . In particular the v independent term p_0 now describes an elliptically fibered Calabi–Yau n -fold Z_n , rather than a torus, and p_+ defines a family of holomorphic stable bundles on Z_n . The fiber construction can be done most easily within the framework of toric geometry as will be described in sect.6.

4. Heterotic/type IIA and heterotic/F-theory duality

We started the discussion of stable holomorphic vector bundles on Calabi–Yau n -folds in the framework of heterotic $\mathcal{N} = 1$ string vacua, and arrived at the same physics in the last section using only type IIA physics. In particular we are free to interpret these type IIA data as valid classical heterotic vacua. We thus arrive at the conclusion that the two theories are the same in the proposed limits - the local mirror limit of type IIA and the perturbative low energy limit of heterotic string. This is the "classical" part of a conjectured duality of full quantum string theories, namely between type IIA on a Calabi–Yau $n + 1$ -fold $\mathcal{W}_{n+1}$ and heterotic string compactified on a $Z_n \times T^2$, with Z_n a Calabi–Yau n -fold. Note that we used only type IIA physics to derive our results and therefore proved such an equivalence on the classical level rather than assuming or using it.

Actually we would like to get rid off the T^2 factor on the heterotic side. The fact that we used elliptically fibered geometries⁵ makes it possible to take what is called the F-theory [7] limit: the limit in which the Kähler class of the elliptic fiber shrinks to zero size⁶. Note that this limit can be taken in $\mathcal{W}_{n+1}$ without interfering with our construction since we concentrated on the complex deformations, rather than the Kähler deformations.

Note that the geometric construction of holomorphic stable bundles on Calabi–Yau n -folds Z_n in terms of local geometries works for any rank of the structure group H . However precisely if H fits into the primordial gauge symmetry of the heterotic string, that is $E_8 \times E_8$ or $SO(32)$, the local geometries $\mathcal{W}_{n+1}$ can be embedded in global compact Calabi–Yau's $\mathcal{W}_{n+1}$. In this case, from the point of F-theory/heterotic string duality, we have obtained a systematic way to construct dual pairs: we associate to the Calabi–Yau $n + 1$ fold $\mathcal{W}_{n+1}$ used for F-theory compactification the heterotic data of a Calabi–Yau n -fold Z_n and a bundle on it (and *vice versa*).

⁵ This requires that the mirror geometries $\mathcal{W}_2$ of the elliptically fibered geometries $\mathcal{M}_2$ are also elliptically fibered. This is the case.

⁶ See also [30].

5. Holomorphic stable bundles on elliptic Calabi–Yau manifolds

In the next section we will sketch the precise formulation of the geometric construction of holomorphic stable bundles on elliptic Calabi–Yau manifolds in terms of toric geometry. Before doing that, we outline some qualitative properties of these bundles and the geometric approach to them.

5.1. Some known results

We proceed with an informal collection of some known facts about holomorphic stable vector bundles on Calabi–Yau manifolds Z . The interested reader can find the details in the literature, as quoted. Most of the concepts in the following sections can be followed without being familiar with the definitions and facts presented in this paragraph.

- A holomorphic stable bundle V on an elliptic curve E is simply a flat bundle on E . It is specified by a representation of the abelian fundamental group of E in a maximal torus of the structure group H (assuming the latter is simply connected).
- [25]: The moduli space of flat H bundles on a fixed E is a weighted projective space $\mathbf{WP}_{s_0, \dots, s_r}^r \equiv W^r$. Here $r = \text{rank } H$ and s_i are the Dynkin indices of the affine Dynkin diagram associated to H (see appendix A.).
- [13]: A geometric realization of the moduli space $\mathcal{M}_E$ of flat bundles on E (including the deformations of E) can be given as follows: for $H = A_{N-1}$, V is specified by N unordered points on the Jacobian $\hat{E}$ of E (corresponding to the values of the Wilson lines in the fundamental representation). If E is given in Weierstrass form

$$p_0 = y^2 + 4x^3 - g_2x - g_3 = 0, \quad (5.1)$$

the N points can be defined as the roots of a degree N polynomial

$$p_C = a_N + a_{N-2}x + a_{N-3}y + \dots + a_0x^{N/2} = 0 \quad (5.2)$$

for N even, the last term being $a_0yx^{(N-3)/2}$ for N odd. The coefficients a_k parametrize the $\mathbf{P}^{N-1}$ predicted in [25].

For $H = E_k$, $k = 6, 7, 8$, a geometric description can be obtained from del Pezzo surfaces dP_k . E.g., for $k = 8$, the equation

$$\begin{aligned} y^2 = & 4x^3 - g_2xv^4 - g_3v^6 + (\alpha_6u^6 + \alpha_5u^5v + \alpha_4u^4v^2 + \alpha^3u^3v^3 + \alpha^2u^2v^4) \\ & + x(\beta_4u^4 + \beta_3u^3v + \beta_2u^2v^2 + \beta_1uv^3), \end{aligned} \quad (5.3)$$

describes a del Pezzo surface obtained from blowing up eight points in $\mathbf{P}^2$. The elliptic curve is identified with the hyperplane $u = 0$. The coefficients α_k, β_k parametrize the $\mathbf{WP}_{1,2,2,3,3,4,4,5,6}^8$ predicted in [25].

- [13]: Holomorphic stable bundles on elliptically fibered Calabi–Yau manifold Z can be described by fibering the data (E, W^r) holomorphically over a complex base B . The non-trivial fibration of the torus E over B builds up the Calabi–Yau n -fold Z_n . Similarly, the projective spaces W^r fit into a holomorphic bundle $\underline{W}^r$ over B . If $s : B \rightarrow \underline{W}^r$ is a section, the homogeneous coordinates $\tilde{a}_i$ of $\underline{W}^r$ pull back to sections

$$\tilde{a}_i \in H^0(B, \mathcal{N}^{s_i} \otimes \mathcal{L}^{-di}), \quad (5.4)$$

where $\mathcal{L}$ is the anti-canonical bundle of B and $\mathcal{N}$ a line bundle on B which is an important characteristic of V and is closely related to the higher Chern classes of V . The d_j are the degrees of the independent Casimir operators of the group H and s_i the Dynkin indices as above. Part of the data of V are defined by choosing a section of $\underline{W}^r$.

The space $\mathcal{Y}$ of sections of $\underline{W}^r$ is the base of the moduli space of holomorphic flat bundles over Z ; the fiber data include additional continuous and discrete data related to certain twistings of V^7 .

A universal abstract construction of V , the so-called parabolic construction, has been given in terms of deformations of unstable bundles. The method is so far restricted to bundles that are invariant under the involution $y \rightarrow -y$ on the fiber and have $c_3(V) = 0$.

- [13]: Chern classes of V have been calculated in the parabolic and spectral cover approach for $H = SU(N)$:

$$c_2(V) = \eta\sigma - \frac{1}{24}c_1(B)^2(N^3 - N) - \frac{1}{8}N\eta(\eta - Nc_1(B)) - \frac{1}{2}\pi_*(\gamma^2). \quad (5.5)$$

Here π denotes the elliptic fibration $\pi : Z \rightarrow B$, σ the class of the section, $\eta = c_1(\mathcal{N})$ and $\gamma = \lambda(N\sigma - \eta + Nc_1(B))$ is a certain class in $H^{1,1}(C, \mathbf{Z})$ which fulfills $\pi_*\gamma = 0$. Here C is the spectral cover described by an equation of the form (5.2) with a_i replaced by $\tilde{a}_i$ as in (5.4) and moreover $\lambda \in \mathbf{Z}/2$.

The third Chern class $c_3(V)$ has been determined in [33][32]:

$$c_3(V) = 2\lambda\sigma\eta(\eta - Nc_1(B)). \quad (5.6)$$

5.2. The geometric approach

The geometric type II string approach improves in various aspects the previous understanding and use of holomorphic stable bundles and heterotic/F-theory duality. We will obtain

- A canonical description of flat bundles V on a torus E in terms of two-complex dimensional type IIA compactification geometries, provided by a map $f : \mathcal{W}_2 \rightarrow (E, V)$. This unifies the results of [13] for A_N and E_k and generalizes to any structure group H .
- More generally we obtain a map $f : \mathcal{W}_{n+1} \rightarrow (Z_n, V)$ that assigns to a local Calabi–Yau $n + 1$ -fold $\mathcal{W}_{n+1}$ an elliptically fibered Calabi–Yau n -fold Z_n together with a family of holomorphic stable vector bundles V on it.
- All objects will be defined within the framework of toric geometry. Differently than in the methods summarized above, singularities in the elliptic fibration of Z_n and sheaf generalizations of V are automatically taken care of in the toric formulation.
- On the practical side, the method allows us to freely engineer bundles (sheafs) on any elliptically fibered Calabi–Yau manifold Z_n . The defining data, such as the structure group H , the manifold Z_n and characteristic classes of V have an extremely simple representation in terms of the toric polyhedron associated to the geometry $\mathcal{W}_n$.
- Interpreting the data (Z_n, V) in terms of the heterotic string, one obtains valid $\mathcal{N} = 1$ vacua that solve (2.2)(2.4). The coincidence of the type IIA physics on $\mathcal{W}_n$ and heterotic string on (Z_n, V) is related to the conjectured non-perturbative type IIA/heterotic and F-theory/heterotic string dualities. We use

⁷ See [31] for a partial identification of these data and refs. [13][20][32] for comments on the data related to additional singularities in the construction.

classical type IIA physics to derive this equivalence, rather than assuming it. This elucidates type IIA/heterotic duality and F-theory/heterotic duality at the classical level.

- Extending the map f in a global context $f : W_{n+1} \leftrightarrow (Z_n, V)$ provides a systematic identification of dual pairs of F-theory/heterotic duality.
- Important data of the bundle V , such as the line bundle $\mathcal{N}$, stability of V as well as non-perturbative five-branes, have a very transparent interpretation in the toric formulation.

6. Toric geometry: A do-it-yourself kit for (non-perturbative) heterotic $\mathcal{N} = 1$ vacua

After outlining the conceptual framework in the previous sections, we want now to describe a concrete recipe, how to define and construct the complex manifold W_{n+1} such that it describes a chosen "heterotic" Calabi–Yau Z_n and a gauge background V on it⁸. We will define the dual pair W_{n+1} and (Z_n, V) within the framework of toric geometry, which for many purposes is the most useful and general description of Calabi–Yau manifolds. We will see that toric geometry is also quite suited to describe vector bundles on Calabi–Yau's⁹.

The toric construction of the local geometries is described in [18] and we will not repeat the details here. However the toric description has some very simple and graphical aspects which we want to sketch here. Actually we can formulate a set of quite simple rules using the language of toric polyhedra. In particular we will see how the physical quantities, such as the structure group H , the "heterotic" manifold Z_n , data of the characteristic classes of V appearing in [13], singularities of the gauge background as well as associated non-perturbative five-brane wrappings are encoded in a "toric polyhedron".

In a nutshell, toric geometry describes a k dimensional Calabi–Yau geometry M_k by a $k+1$ dimensional polyhedron Δ_{k+1}^* , defined as the convex hull of a set $\{\nu_i^*\}$ of r integral vertices ν_i^* defined in a standard $k+1$ dimensional lattice. Each vertex ν_i^* on a face of Δ_{k+1}^* of codimension larger than one defines a divisor (holomorphic hypersurface) in M_k . Note that because of the dimension of Δ_{k+1}^* , there must be $r - k - 1$ linear relations $l_i^{(m)}$ between the r (external) vertices ν_i^*

$$\sum_i l_i^{(m)} \nu_i^* = 0, \quad m = 1, \dots, h^{1,1}. \quad (6.1)$$

In fact $h^{1,1} = r - k - 1$ is the number of Kähler moduli of M_k .

Now choose the elliptically fibered "heterotic" Calabi–Yau n -fold Z_n with base $\pi : Z_n \rightarrow B_{n-1}$ of the fibration. To Z_n corresponds an $n+1$ dimensional polyhedron $\Delta_{Z_n}^*$. The guideline to construct families of holomorphic stable bundles on Z_n in terms of toric geometry is then as follows [18]:

1. The structure group H :

Choose the structure group H . The polyhedron Δ_3^* for the geometry $\mathcal{W}_2(H)$ and, if existing, a global embedding $\mathcal{W}_2(H)$ is given in [23][18]. The perturbative gauge symmetry of the heterotic string is the commutant of H in $E_8 \times E_8$.

⁸ Since we consider structure groups H which are embeddable in the heterotic gauge group we will treat the local version $\mathcal{W}_{n+1}$ and the global embedding W_{n+1} on the same footing in the following.

⁹ A fact that has been noticed already in the formulation of $(0, 2)$ linear sigma models [10].

2. The base geometry:

The toric base of the elliptic fibration B_{n-1} is described by a toric polyhedron $\Delta_{B_{n-1}}^*$ as given e.g. in [18].

3. The fibration $W_2 \rightarrow W_{n+1} \rightarrow B_{n-1}$:

The two polyhedra are now embedded in a $3 + n - 1$ dimensional polyhedron Δ_{n+2}^* that describes a Calabi–Yau W_{n+1} :

$$\Delta_{W_{n+1}}^* \subset \begin{pmatrix} 0 & \Delta_3^* \\ \Delta_{B_{n-1}}^* & * \end{pmatrix}. \quad (6.2)$$

4. Embedding of Z_n :

To the heterotic compactification manifold Z_n corresponds a $n + 1$ dimensional polyhedron $\Delta_{Z_n}^*$. It appears as a *projection* in $\Delta_{W_{n+1}}^*$. To define the projection we note that the fact that we use elliptically fibered geometries implies [34] that Δ_3^* is itself of a specific form. In our case Δ_3^* splits as

$$\Delta_3^* = \begin{pmatrix} 0 & \Delta_2^* \\ \pm 1 & * \end{pmatrix}, \quad (6.3)$$

where Δ_2^* is the two-dimensional polyhedron defining the elliptic fiber. The projection to the heterotic manifold is defined in the direction transverse to the hyperplane of the elliptic fiber:

$$\begin{pmatrix} 0 & \begin{matrix} \downarrow \\ \Delta_{B_{n-1}} \end{matrix} & \begin{matrix} \Delta_2^* \\ * \end{matrix} \\ \Delta_{B_{n-1}} & * & \end{pmatrix} \quad \Delta_{Z_n}$$

This concludes the description of the geometrical data of the heterotic compactification¹⁰.

5. Perturbative bundle data: fixing the line bundle $\mathcal{N}$ on B_{n-1} :

The above conditions do not fix $\Delta_{W_{n+1}}^*$ completely: there are many $n + 2$ dimensional vertices that project to the $n + 1$ dimensional vertices of $\Delta_{Z_n}^*$ in the above construction. This freedom is related to perturbative and non-perturbative data associated to the gauge background V . Let us first consider the perturbative data. According to [13], the definition of V includes the specification of a line bundle $\mathcal{N}$ on B_{n-1} . The class $\eta = c_1(\mathcal{N})$ enters the Chern classes $c_2(V)$ and $c_3(V)$ as in (5.5)(5.6).

In the toric framework, line bundles are related to the linear relations (6.1) between the vertices of $\Delta_{W_{n+1}}^*$. Recall also that $\Delta_{W_{n+1}}^*$ is required to be convex. Making a minimal choice of vertices $\{\nu_i^*\}$, such that $\Delta_{W_{n+1}}^*$ fulfils the conditions 1.-4., and moreover is convex, determines obviously the linear relations (6.1) of the vertices in $\Delta_{W_{n+1}}^*$ and thus in particular the line bundles on B_{n-1} . Specifically it turns out that there are only two line bundles that determine the fibration of the two complex-dimensional geometries W_2 over B_{n-1} , namely the anti-canonical bundle $\mathcal{L}$ of B_{n-1} and one additional line bundle which is equivalent to $\mathcal{N}$ of ref.[13].

Thus determining the bundle data $\mathcal{N}$, translates in the toric polyhedra to a simple linear relation

¹⁰ Note that there can be obstructions to give all curves in Z_n a finite size; this phenomenon is related to non-perturbative gauge symmetries [18].

of the vertices in $\Delta_{W_{n+1}}^*$. In fact the definition in toric geometry is more general in the sense that it applies to any singularities in the geometry or the gauge background. An interesting fact is that the convexity of $\Delta_{W_{n+1}}^*$ restricts the possible linear relations and thus the range of acceptable line bundles $\mathcal{N}$. In six dimensions this phenomenon describes the well-known stability properties of instantons on K3 and we expect a similar interpretation in four dimensions. A general bound on the first Chern class $\eta = c_1(\mathcal{N})$ is given by

$$\nu(G) c_1(\mathcal{L}) \leq \eta \quad (\leq 12c_1(\mathcal{L})), \quad (6.4)$$

where G is the singularity type of the fiber geometry W_2 and $\nu(G)$ is a certain characteristic number defined in [35]. Explicit expressions $\mathcal{N}$ for toric bases, such as $\mathbf{P}^2$ and $\mathbf{F}_n$ and blow ups thereof, can also be found in [18]. See also ref.[36] for an independent discussion of the class η in terms of toric polyhedra¹¹.

6. Non-perturbative bundle data I: gauge symmetries :

After fixing H , Z_n and the line bundle $\mathcal{N}$, there is still a freedom to add vertices to $\Delta_{W_{n+1}}^*$. This freedom corresponds to introducing singularities in the behavior of V that are related to non-perturbative physics of the heterotic string. Specifically we can introduce singularities in the elliptic fibration of W_{n+1} located over a divisor of B_{n-1} . Gauge symmetry enhancement follows from the physics of the type IIA/F-theory compactification on W_{n+1} [8]. In the heterotic picture these gauge symmetries are non-perturbative. Amazingly, the simple requirement of convexity of $\Delta_{W_{n+1}}^*$ gives an almost complete information about the possible complicated non-perturbative gauge symmetries, e.g. in the case of five-branes at singularities [12]. We will describe some singularities of the heterotic bundle associated to this non-perturbative gauge enhancements in the next section.

7. Non-perturbative bundle data II: non-perturbative five-branes :

We can also blow up the base $\tilde{B}_n$ of the *elliptic* fibration of W_{n+1} . It is a $\mathbf{P}^1$ bundle over the heterotic base B_{n-1} . In the six dimensional case, corresponding to $n = 1$ and a K3 compactification of the heterotic string, blow ups of the $\mathbf{P}^1$ fibration have been identified as non-perturbative heterotic 5 branes filling space time and located at points of the K3 [8]. The story in four dimensions is similar [18], with the difference that the five-branes are now wrapped on holomorphic spheres C in B_{n-1} specified by the position of the blow-up¹². For C a genus zero curve the universal spectrum from the five-branes is an $\mathcal{N} = 1$ tensor multiplet L . It couples to a non-critical string in four dimensions. This is a quite interesting object from the point of supersymmetric $\mathcal{N} = 1$ particle physics. In particular the string tension can be classically zero but acquires a small, non-perturbatively generated tension $\sim \Lambda^{-2}$ at the quantum level [37], very similarly as the non-perturbative mass of the $W^\pm$ bosons in the $\mathcal{N} = 2$ $SU(2)$ Seiberg–Witten model. It would be interesting to study their implication for standard particle physics below the scale Λ .

This concludes our recipe for the construction of the heterotic $\mathcal{N} = 1$ vacuum. The precise information about which toric vertices implement the points 1.-7. can be found in [18]. It is striking how simple toric concepts, such as convexity of $\Delta_{W_{n+1}}^*$, contain interesting physics. Moreover singularities, both of the geometry Z_n and the bundle V can be easily introduced in the toric language. However we should also mention that there

¹¹ The bound on η derived in [36] is different from (6.4).

¹² See also ref. [36] for a discussion of these five-branes.

are few additional data of V [13] which are not encoded in the simple toric description above. It would be interesting to have a generalization that contains them.

Note that apart from the classical geometric formulation of (families of) holomorphic stable gauge backgrounds on Z_n , we can think of the correspondence $W_{n+1} \leftrightarrow (Z_n, V)$ as a dual pair and assume, use - or possibly verify - duality of the *quantum* theories, to study quantum and non-perturbative phenomena (as we did in fact already, when we used F-theory to infer non-perturbative heterotic phenomena). Some interesting directions that have been considered include correlation functions and the $\mathcal{N} = 1$ superpotential. See refs. [38] for examples.

7. Some results

Let us finally sketch some results on (non-perturbative) heterotic physics which have been obtained in [18] using the above method.

7.1. Standard embedding

A simple solution to the equations (2.1),(2.2) is to set the gauge connection equal to the spin connection of the manifold, $V = TZ$. Though this configuration is not too appealing for phenomenology - the heterotic gauge group is $E_6 \times E_8$ and the matter spectrum is tied to the Euler number of Z - it has been studied extensively in the past because it was one of the few known solutions. The structure group of V is then $SU(n)$ for a Calabi–Yau n -fold, and the local geometry described by an equation of the form (3.7).

The global geometry W_{n+1} corresponding to the F-theory dual can be determined in the following way. From the fact that the bundle in the second factor is trivial it follows that the line bundle $\mathcal{N}$ discussed in sect.6 is trivial in the second E_8 factor. The Weierstrass form for W_{n+1} and Z_n takes the following form:

$$\begin{aligned} p_{Z_n} &= y^2 + x^3 + x\tilde{z}^4 f + \tilde{z}^6 g, \\ p_{W_n} &= y^2 + x^3 + x(\tilde{z}zw)^4 f + (\tilde{z}zw)^6 g + \tilde{z}^6 z^5 w^7 \Delta + \tilde{z}^6 z^7 w^5. \end{aligned} \tag{7.1}$$

Here (z, w) denote the variables parametrizing the base $\mathbf{P}^1$ of the elliptically fibered K3 fiber W_2 of the fibration $\pi_F : W_{n+1} \rightarrow B_{n-1}$ and $\Delta = 4f^3 + 27g^2$ is the discriminant of the elliptic fibration of Z_n . This generalizes the six-dimensional result obtained in [39] to four and lower dimensions.

7.2. Non-perturbative gauge symmetries

In the F-theory picture gauge symmetries correspond to singularities in the elliptic fibration. Those located over the base B_{n-1} have a perturbative interpretation in the heterotic dual, whereas non-perturbative ones are located over curves in B_{n-1} [8]. Using the map $W_{n+1} \rightarrow (Z_n, V)$ we can associate the heterotic compactification that creates the same dynamics non-perturbatively. In the six-dimensional case we find:

(*) *Consider the $E_8 \times E_8$ string compactified on an elliptically fibered K3 with a singularity of type G at a point $s = 0$ and a special gauge background $\hat{V}$. If the restriction of the spectral cover¹³ of $\hat{V}|_E$ to the fiber E at $s = 0$ is sufficiently trivial, the heterotic string acquires a non-perturbative gauge symmetry $G_{np} \supset G$.*

The triviality condition can be made precise by specifying the behavior of V near $s = 0$ [18]. Similar results hold for four-dimensional compactifications.

We point out that the above triviality condition on the spectral cover does not imply that the field strength of V , which measures the behavior of V near the singularity, is trivial. In fact it has been shown recently in [40] that if $F = 0$ on the singularity, the conformal field theory of the heterotic string is well behaved and there is no non-perturbative gauge symmetry.

7.3. Non-perturbative dualities

Our map $f : W_{n+1} \rightarrow (Z_n, V)$ can be ambiguous in the sense that there are two (or even more) ways to associate a pair (Z_n, V) to $\Delta_{W_{n+1}}^*$. If the two maps are compatible with the same elliptic fibration, we obtain a non-perturbative duality of two heterotic string theories

$$(Z_n, V) \sim (Z'_n, V'). \quad (7.2)$$

The conditions under which this duality exists, can be formulated in simple properties of the polyhedron $\Delta_{W_{n+1}}^*$ associated to W_{n+1} (essentially the existence of appropriate hyperplanes and projections in $\Delta_{W_{n+1}}^*$) [18]. The generic form of the duality is the following:

(**) *Let the heterotic string be compactified on a Calabi–Yau three-fold with G' singularity and with a certain gauge background with structure group H such that the toric data $\Delta_{W_{n+1}}^*$ fulfil the above mentioned condition. Then there exists a non-perturbatively equivalent compactification on a Calabi–Yau manifold with G singularity and with a specific gauge background with structure group H' .*

Here H (H') is the commutant of G (G') in $E_8 \times E_8$. Note that the duality exchanges the groups of the geometric singularity and the gauge bundle: the singularity in the dual manifold is the commutant of the structure group and *vice versa*. Let us give an extreme example of such an duality: the heterotic string with a trivial gauge bundle on a smooth K3 has a perturbative $E_8 \times E_8$ gauge symmetry and in addition $n'_T = 24$ non-perturbative tensor multiplets from the 24 five-branes required to satisfy (2.5). The dual theory is a heterotic theory with $E_8 \times E_8$ gauge bundle on a K3 with $E_8 \times E_8$ singularity. The perturbative gauge symmetry is trivial, while non-perturbative dynamics associated to the singularity produce both, the $E_8 \times E_8$ gauge symmetry as well as 24 extra tensor multiplets.

This duality reminds very much of a known duality in the linear sigma model formulation of heterotic strings, namely a symmetry of the formulation under the exchange of the data defining the manifold and the data defining the bundle [41]. However note that in our case this duality is in general *non-perturbative*.

¹³ For $G \neq SU(N)$ we use the generalizations of the spectral cover discussed previously.

7.4. Mirror symmetry of F-theory

Consider a six-dimensional F-theory compactification on W_3 and the associated heterotic dual (Z_2, V) as defined above. If the mirror manifold M_3 of W_3 is also elliptically fibered, there is also a heterotic theory (Z'_2, V') associated to it and one can ask the question of how the two theories are related. The answer is that after compactification on a three-torus to three dimensions, the two become equivalent in virtue of mirror symmetry of type II strings [42].

A proposal for the relation between (Z_2, V) and (Z'_2, V') has been made in [43] based on a comparison of hodge numbers and gauge symmetries in some cases. Using the toric map $W_{n+1} \rightarrow (Z_n, V)$ one can derive the two theories; in fact one finds a subtle realization of Higgs and Coulomb branches in toric geometry as expected from the action of three-dimensional mirror symmetry [18]. The general relation is of a similar form as in (**) above: a heterotic theory with a G bundle compactified on a K3 manifold with $\hat{H}$ singularity gets mapped to a heterotic theory with a bundle with structure group $\hat{H}$ compactified on a manifold with G singularity. Note that we have an explicit map of the moduli spaces of the two theories. It would be interesting to formulate the associated three-dimensional dual pairs associated to this relation, as in [44].

8. Outlook

In these lectures we have reviewed a construction of a certain class of Calabi–Yau geometries W_{n+1} and, if possible, their compact embeddings W_{n+1} . In a certain limit of the type IIA string compactified on W_{n+1} we obtain the physics of holomorphic, stable vector bundles (or sheaves) V on a Calabi–Yau manifold Z_n . Interpreting the data (Z_n, V) as a valid classic heterotic vacuum establishes F-theory/heterotic duality at a classical level. In the toric construction, we have a well-defined toric map $f : W_{n+1} \rightarrow (V, Z_n)$ that allows for a systematic construction of dual pairs in both directions.

The identifications between W_{n+1} and (Z_n, V) may serve as a starting point to study more refined non-perturbative relations between the dual F- and heterotic string theory. In particular we would like to formulate both, perturbative and non-perturbative quantities of the heterotic string, e.g. gauge and Yukawa couplings of the $(0, 2)$ vacua, in terms of geometric quantities on W_{n+1} , such as the correlation functions determined by period integrals considered in [37]. In the ideal (and supposedly too optimistic) case, we can hope to obtain exact results for at least some of the holomorphic physical quantities in the $\mathcal{N} = 1$ theory from geometric information on W_{n+1} , similarly as it happened to work in the case of $\mathcal{N} = 2$ supersymmetry [6][45][24].

Acknowledgements:


The author would like to thank the organizers for the opportunity to present this lectures and their hospitality and P. Berglund, S. Katz and C. Vafa for the pleasant collaboration on the subjects presented above.

Appendix A. Local mirror geometries $\mathcal{W}_2$

On the right, the affine Dynkin diagrams together with the Dynkin indices s_i are shown. The polynomial $p_{\mathcal{W}_2} = 0$ for the associated local complex geometry $\mathcal{W}_2$ is shown on the left. Each node in the Dynkin diagram of index s_i contributes precisely one monomial with v power s_i to $p_{\mathcal{W}_2}$.

$$\mathbf{A}_{N-1} : \mathbf{v}^0 (y^2 + x^3 + z^6 + yxz) +$$


$$\mathbf{v}^1 (z^N + z^{N-2}x + z^{N-3}y + \dots + \left\{ \begin{matrix} x^{N/2} \\ yx^{\frac{N-3}{2}} \end{matrix} \right\})$$



$$\mathbf{B}_N : \mathbf{v}^0 (y^2 + x^3 + z^6 + xyz) +$$

$$\mathbf{v} (z^N + z^{N-3}y + \epsilon yx^{(N+1)/2-2} + (1-\epsilon)x^{N/2}) +$$

$$\mathbf{v}^2 (z^{2N-6} + z^{2N-8}x + z^{2N-10}x^2 + \dots + x^{N-3})$$



$$\mathbf{C}_N : \mathbf{v}^0 (y^2 + x^3 + z^6 + yxz) +$$


$$\mathbf{v} (z^{2N} + z^{2N-2}x + z^{2N-4}x^2 + \dots + x^N)$$



$$\mathbf{D}_N : \mathbf{v}^0 (y^2 + x^3 + z^6 + xyz) +$$


$$\mathbf{v} (z^{N-3}y + z^N + z^{1-\epsilon}yx^{(N+\epsilon)/2-2} + z^\epsilon x^{(N-\epsilon)/2}) +$$

$$\mathbf{v}^2 (z^{2N-6} + z^{2N-8}x + z^{2N-10}x^2 + \dots + z^2x^{N-4})$$




$$\mathbf{G}_2 : \mathbf{v}^0 (y^2 + x^3 + z^6 + yxz) +$$

$$\mathbf{v} (z^3 + y) + \mathbf{v}^2$$




$$\mathbf{F}_4 : \mathbf{v}^0 (y^2 + x^3 + z^6 + yxz) +$$

$$\mathbf{v} (z^4 + x^2) + \mathbf{v}^2 (z^2 + x) + \mathbf{v}^3$$




$$\mathbf{E}_6 : \mathbf{v}^0 (y^2 + x^3 + z^6 + xyz) +$$

$$\mathbf{v} (z^5 + zx^2 + xy) + \mathbf{v}^2 (z^4 + z^2x + zy) + \mathbf{v}^3 z^3$$



$$\mathbf{E}_7 : \mathbf{v}^0 (y^2 + x^3 + z^6 + xyz) +$$


$$\mathbf{v} (z^5 + xy) + \mathbf{v}^2 (z^4 + zy + x^2) + \mathbf{v}^3 (z^3 + zx) + \mathbf{v}^4 z^2$$



$$\mathbf{E}_8 : \mathbf{v}^0 (y^2 + x^3 + z^6 + xyz) +$$

$$\mathbf{v} (z^5) + \mathbf{v}^2 (z^4 + x^2) + \mathbf{v}^3 (z^3 + y) +$$

$$\mathbf{v}^4 (z^2 + x) + \mathbf{v}^5 z + \mathbf{v}^6$$



References

- [1] P. Candelas, G.T. Horowitz, A. Strominger and E. Witten, Nucl. Phys. B258 (1985) 46
- [2] C. Hull and E. Witten, Phys. Lett. B160 (1985) 398.;
E. Witten, Nucl. Phys. B268 (1986) 79.
- [3] J. Distler, Phys. Lett. B188 (1987) 431; J. Distler and B. Greene, Nucl. Phys. B304 (1988) 1.
- [4] C.M. Hull and P.K. Townsend, Nucl. Phys. B438 (1995) 109
- [5] E. Witten, Nucl. Phys. B443 (1995) 85
- [6] S. Kachru and C. Vafa, Nucl. Phys. B450 (1995) 69;
S. Ferrara, J.A. Harvey, A. Strominger and C. Vafa, Phys. Lett. B361 (1995) 59.
- [7] C. Vafa, Nucl. Phys. B469 (1996) 403
- [8] C. Vafa and D. Morrison, Nucl. Phys. B473 (1996) 74; Nucl. Phys. B476 (1996) 437.
- [9] M.J. Duff, R. Minasian and E. Witten, Nucl. Phys. B465 (1996) 413.
- [10] J. Distler, B.R. Greene and D. Morrison, Nucl. Phys. B481 (1996) 289.
- [11] O. Ganor and A. Hanany, Nucl. Phys. B474 (1996) 122;
N. Seiberg and E. Witten, Nucl. Phys. B471 (1996) 121.
- [12] J.D. Blum and K. Intriligator, Nucl. Phys. B506 (1997) 199; Nucl. Phys. B506 (1997) 223;
P.S. Aspinwall and D.R. Morrison, Nucl. Phys. B503 (1997) 533;
P. Candelas, E. Perevalov and G. Rajesh Nucl. Phys. B507 (1997) 445.
- [13] R. Friedman, J.W. Morgan and E. Witten, Comm. Math. Phys. 187 (1997) 679.
- [14] E. Witten, Nucl. Phys. B460 (1996) 541.
- [15] K. Becker, M. Becker and A. Strominger, Nucl. Phys. B456 (1995) 130
- [16] M. Bershadsky, V. Sadov and C. Vafa, Nucl. Phys. B463 (1996) 420.
- [17] A. Klemm, W. Lerche, P. Mayr, C. Vafa, N. Warner, Nucl. Phys. B477 (1996) 746.
- [18] P. Berglund and P. Mayr, *Heterotic string/F-theory duality from mirror symmetry*, hep-th/9811217.
- [19] K. Uhlenbeck and S.T. Yau, *On the existence of hermitian Yang-Mills connections in stable vector bundles*, preprint (1996).
- [20] M. Bershadsky, A. Johansen, T. Pantev and V. Sadov, Nucl. Phys. B505 (1997) 165.
- [21] R.Y. Donagi, Asian J. Math. 1 (1997) 214; MSRI pub. 28 (1992) 65.
- [22] C. Vafa and E. Witten, Nucl. Phys. Proc. Suppl. 46 (1996) 225.
- [23] S. Katz, P. Mayr and C. Vafa, Adv. Theor. Math. Phys. 1 (1998) 53.
- [24] A. Klemm, W. Lerche, P. Mayr, C. Vafa, N. Warner, Nucl. Phys. B477 (1996) 746;
S. Katz, A. Klemm and C. Vafa, Nucl. Phys. B497 (1997) 173;
S. Katz, P. Mayr and C. Vafa, Adv. Theor. Math. Phys. 1 (1998) 53.
- [25] E. Looijenga, Invent. Math. 38 (1977) 17; Invent. Math. 61 (1980) 1.
- [26] E. Witten, J. High Energy Phys. 02 (1998) 6.
- [27] S. Chaudhuri, G. Hockney and J. D. Lykken, Phys. Rev. Lett. 75 (1995) 2264.

- [28] P. Berglund, A. Klemm, P. Mayr and S. Theisen, *On type IIB vacua with varying coupling constant*, hep-th/9805189.
- [29] C. Vafa and E. Witten, Nucl. Phys. Proc. Suppl. 46 (1996) 225.
- [30] P. Aspinwall, *K3 surfaces and string duality*, hep-th/9611137
- [31] G. Curio and R.Y. Donagi, Nucl. Phys. B518 (1998) 603.
- [32] G. Curio, Phys. Lett. B435 (1998) 39.
- [33] B. Andreas, *On vector bundles and chiral matter in $N=1$ heterotic compactifications*, hep-th/9802202
- [34] A. C. Avram, M. Kreuzer, M. Mandelberg and H. Skarke, Nucl. Phys. B494 (1997) 567.
- [35] P. Berglund and P. Mayr, *Stability of vector bundles from F -theory*, hep-th/9904114.
- [36] G. Rajesh, J. High Energy Phys. 12 (1998) 18.
- [37] P. Mayr, Nucl. Phys. B494 (1997) 489.
- [38] E. Witten, Nucl. Phys. B474 (1996) 343; P. Mayr, Nucl. Phys. B494 (1997) 489; S. Katz and C. Vafa, Nucl. Phys. B497 (1997) 204; C. Vafa, Adv.Theor.Math.Phys. 2 (1998) 497; W. Lerche, J. High Energy Phys. 11 (1997) 4; W. Lerche and S. Stieberger, *Prepotential, mirror map and F theory on $K3$* , hep-th/9804176; W. Lerche, S. Stieberger and N.P. Warner, hep-th/9811228; hep-th/9901162.
- [39] P.S. Aspinwall and R.Y. Donagi, *The Heterotic string, the tangent bundle, and derived categories*, hep-th/9806094.
- [40] E. Witten, *Heterotic string conformal field theory and A - D - E singularities*, hep-th/9909229.
- [41] J. Distler and S. Kachru, Nucl. Phys. B442 (1995) 64.
- [42] K. Intriligator and N. Seiberg, Phys. Lett. B387 (1996) 513.
- [43] E. Perevalov and G. Rajesh, Phys. Rev. Lett. 79 (1997) 2931.
- [44] K. Hori, H. Ooguri and C. Vafa, Nucl. Phys. B504 (1997) 147.
- [45] S. Kachru, A. Klemm, W. Lerche, P. Mayr and C. Vafa, Nucl. Phys. B459 (1996) 537.